




Generating Functions

- Readings:
Rosen section 6.4



Generating Functions

- Represent sequences $\{a_n\}$ by coding the term a_n as the coefficient of x^n in a power series.
 $a_n = 1, 2, 4, 8, \dots \rightarrow G(x) = 1 + 2x + 4x^2 + \dots$
 $b_n = 1, -1, 1, -1, \dots \rightarrow G(x) = 1 - x + x^2 - \dots$
- The generating function for the sequence $a_n = a_0, a_1, a_2, \dots$ of real numbers is the infinite series

$$G(x) = a_0 + a_1x + a_2x^2 + \dots = \sum_{k=0}^{\infty} a_k x^k$$



- Example (Rosen p.436):
Find the generating functions for the sequences $\{a_k\}$ with:

$$a_k = 3$$

$$a_k = k + 1$$

$$a_k = 2^k$$



Useful Facts about Power Series

- $1/(1-x) = 1 + x + x^2 + \dots$ for $|x| < 1$
- $1/(1-ax) = 1 + ax + ax^2 + \dots$ for $|ax| < 1$

Adding & multiplying two generating functions

$$\text{Let } f(x) = \sum_{k=0}^{\infty} a_k x^k \text{ and } g(x) = \sum_{k=0}^{\infty} b_k x^k$$

$$f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$$

$$f(x)g(x) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k a_j b_{k-j} \right) x^k$$



- Example (Rosen p.437):

Let $f(x) = \frac{1}{(1-x)^2}$. Find the coefficients in the

expansion $f(x) = \sum_{k=0}^{\infty} a_k x^k$



Extended Binomial Coefficient

- To apply binomial theorem for exponents that are not positive integers.

Let u be a real number and k a nonnegative integer. Then the extended binomial coefficient, $\binom{u}{k}$, is defined by:

$$\binom{u}{k} = \begin{cases} u(u-1)\dots(u-k+1)/k! & \text{if } k > 0 \\ 1 & \text{if } k = 0 \end{cases}$$



- Example (Rosen p.438):

$$\binom{-2}{3} =$$

$$\binom{1/2}{3} =$$



- When the top parameter is a negative number, the extended binomial coefficient can be expressed in terms of an ordinary binomial coefficient.

$$\binom{-n}{r} =$$



Extended Binomial Theorem

- Let x be a real number with $|x| < 1$ and let u be a real number. Then

$$(1+x)^u = \sum_{k=0}^{\infty} \binom{u}{k} x^k$$

$$(1+x)^{-n} =$$

$$(1-x)^{-n} =$$



Counting Problems and Generating Functions

Example (Rosen p.441):

Find the number of solutions of $e_1 + e_2 + e_3 = 17$ where $2 \leq e_1 \leq 5$, $3 \leq e_2 \leq 6$, $4 \leq e_3 \leq 7$



- Example (Rosen p.441):

Find the number of ways to insert tokens worth \$1, \$2, and \$5 into a vending machine to pay for an item that costs \$r:

When order **does not** matter.



- Example (Rosen p.441):

Find the number of ways to insert tokens worth \$1, \$2, and \$5 into a vending machine to pay for an item that costs \$r:

When order **does** matter.



- Example (Rosen p.442):

Use generating functions to find the number of *k-combination* of a set with n elements. (Assume that the binomial theorem has been established.)



- Example (Rosen p.443):

Use generating functions to find the number of *r-combination* of a set with n elements when repetition of elements is allowed.



- Example (Rosen p.443):

Use generating functions to find the number of ways to selected r objects of n different kinds if we ***must select at least one object of each kind.***



Using Generating Functions to Solve Recurrence Relations

- Example (p.444):

$$a_k = 3a_{k-1} \text{ for } k=1,2,3,\dots \text{ and } a_0=2$$



- Example (Rosen p.445):
 $a_n = 8a_{n-1} + 10^{n-1}$ and $a_0=1$

Proving Identities via Generating Functions



- Example (Rosen p.446):
Use generating functions to show that

$$\sum_{k=0}^n c(n,k)^2 = c(2n,n)$$